

Open Research Online

The Open University's repository of research publications and other research outputs

Hay Days: management of floodplain meadows for sustainable hay production

Other

How to cite:

Bowskill, Vicky (2021). Hay Days: management of floodplain meadows for sustainable hay production. Postgraduate Research Poster Competition, The Open University.

For guidance on citations see [FAQs](#).

© 2021 Victoria Bowskill



<https://creativecommons.org/licenses/by/4.0/>

Version: Poster

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk

Hay Days: management of floodplain meadows for sustainable hay production

PhD researcher: Vicky Bowskill, Supervisors: David Gowing and Shonil Bhagwat
School of Environment, Earth and Ecosystem Sciences




Floodplain meadows

These internationally threatened meadows depend on annual hay cutting and aftermath grazing to maintain their characteristic botanical diversity. Changes in agricultural methods and land use over the last century have led to the loss of 97% of our heritage wildflower meadows. This study investigates how agricultural production can be brought back into balance with biodiversity conservation through well-timed haymaking.



Hay Days




www.vickybowskill.com

The floodplain meadow nutrient pump

Plant growth: nutrients transferred

Plant growth transfers soil nutrients to biomass during growth and flowering. Nutrients are reabsorbed into underground reserves after flowering.




Haymaking: nutrients removed

Haymaking before flowering finishes removes soil nutrients, preventing them from building up in the soil and ensuring a wide range of plant species can thrive.




Flood sediments: nutrients deposited

Flooding deposits nutrient-rich sediments that enrich soils. Rising soil nutrients can lead to falling botanical diversity as tall competitive species crowd out lower growing plants.




Meadow Managers

Meadows would not be meadows without meadow managers. So this project began by seeking their views to inform the fieldwork focus.



The questionnaire



The interviews



Nature Friendly Farming


Conservation and farming are often portrayed as opposing forces in the media and it became clear from the questionnaire that this divide is inaccurate and unhelpful.



Hay yield and minerals

Floodplain meadows are a naturally fertile farm resource.

How hay makes meadows




Cutting date

Peak yield and peak nutrient content do not occur at the same time, so land managers need to find the right balance. Mineral data for this study is currently being analysed.

Agri-environment schemes often restrict hay cutting dates until after mid-July and this can cause conflict between agricultural and biodiversity management goals.


Dry matter yield on floodplain meadows



Double cutting

One method to maximise yield and total nutrient removal is to take a double hay cut. The aim is to take two harvests during the active growth periods in the early summer and autumn, avoiding the summer dormant period.

Growth and yield on floodplain meadows



Acknowledgements and References

McGinlay, J., Gowing, D. J. G., & Budds, J. (2017). The threat of abandonment in socio-ecological landscapes: Farmers' motivations and perspectives on high nature value grassland conservation. *Environmental Science and Policy*, 69, 39–49. <https://doi.org/10.1016/j.envsci.2016.12.007>
Pavlu, K., Kassahun, T., Pavlu, V. V., Pavlu, L., Blazek, P., & Homolka, P. (2021). The effects of first defoliation and previous management intensity on forage quality of a semi-natural species-rich grassland. *PLoS ONE*, 16(3). <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0248804>
Rothero, E.; Lake, S.; Gowing, D. (2016). Floodplain Meadows - Beauty and Utility A Technical Handbook. In *Floodplain Meadows - Beauty and Utility A Technical Handbook*. http://www.floodplainmeadows.org.uk/sites/www.floodplainmeadows.org.uk/files/Floodplain_Meadows_-_Beauty_and_UTILITY_A_Technical_Handbook.pdf
Schlegel, P., Wyss, U., Arrigo, Y., & Hess, H. D. (2016). Mineral concentrations of fresh herbage from mixed grassland as influenced by botanical composition, harvest time and growth stage. *Animal Feed Science and Technology*, 219, 226–233. <https://doi.org/10.1016/j.anifeeds.2016.06.022>

All illustrations, infographics and photographs on this poster are original work ©Vicky Bowskill. This study runs from 2019 to 2023 with thanks my supervisors, the Floodplain Meadows Partnership, the Open University School of Environment, Earth and Ecosystem Sciences, funder CENTA and study site owners BBOWT, FAI Farms Ltd and The Parks Trust.

